

Physics potential of a few hundred GeV $\mu^+\mu^-$ collider

David B. Cline

University of California Los Angeles, Center for Advanced Accelerators, Physics Department, 405 Hilgard Ave., Los Angeles, CA 90024–1547, USA

Received 8 April 1994

There is growing evidence that both a Standard Model and a SUSY Model Higgs should exhibit one resonance at a mass less than $2M_Z$. This is precisely in the mass range that is very difficult (but not impossible) to detect at the LHC and possibly beyond the reach of LEP II. At a $\mu\mu$ collider the direct channel $\mu^+\mu^- \rightarrow h^0 \rightarrow b\bar{b}$ can be used to search for the Higgs. We discuss the collider requirements for this search on the $\mu\mu$ collider luminosity and the detector. These results arose from a $\mu^+\mu^-$ Collider Workshop held in Napa Valley, California, November 1992.

1. Introduction

At the Port Jefferson Advanced Accelerator Workshop in the Summer of 1992, a group investigating new concepts of colliders studied anew the possibility of a $\mu^+\mu^-$ collider since e^+e^- colliders will be very difficult in the several TeV range [1]. A small group also discussed the possibility of a $\mu^+\mu^-$ collider. A special workshop was then held in Napa, California in the fall of 1992 for this study. There are new accelerator possibilities for the development of such a machine, possibly at an existing or soon to exist storage ring [2, 3]. For the purpose of the discussion here, a $\mu^+\mu^-$ collider is schematically shown in Fig. 1. In this brief note we study one of the most interesting goals of a $\mu^+\mu^-$ collider: the discovery of a Higgs boson in the mass range beyond that to be covered by LEP I&II (~ 80 – 90 GeV) and the natural range of the supercolliders $\geq 2M_Z$ [4]. In this mass range, as far as we know, the dominant decay mode of the h^0 will be

$$h^0 \rightarrow b\bar{b}, \quad (1)$$

whereas the Higgs will be produced by the direct channel

$$\mu^+\mu^- \rightarrow h^0 \quad (2)$$

which has a cross section enhanced by

$$(M_\mu/M_c)^2 \sim (200)^2 = 4 \times 10^4 \quad (3)$$

larger than the corresponding direct product at an e^+e^- collider. However, we will see that the narrow width of the Higgs partially reduces this enhancement.

There is growing evidence that the Higgs should exist in this low mass range from

- 1) the original paper of Cabibbo et al., which shows

that, when $M_t > M_Z$ and assuming a Grand Unification of Forces, $M_h < 2M_Z$ [4] (Fig. 2),

2) fits to LEP data, which imply a low mass h^0 could be consistent with $M_t > 150$ GeV [4],

3) the extrapolation to the GUT Scale that is consistent with SUSY also implies that one of the Higgs should have a low mass, perhaps below 130–150 GeV [4].

This evidence implies the exciting possibility that the Higgs mass is just beyond the reach of LEP II and in a range that is very difficult for the super-colliders to extract the signal from background, ie. either $h^0 \rightarrow \gamma\gamma$ or the very rare $h^0 \rightarrow \mu\mu\mu\mu$ in this mass range, since $h \rightarrow b\bar{b}$ is swamped by hadronic background. However, detectors for the LHC are designed to extract this signal [5].

In this low mass region the Higgs is also expected to be a fairly narrow resonance and thus the signal should stand out clearly from the background from

$$\begin{aligned} \mu^+\mu^- \rightarrow \gamma \rightarrow b\bar{b} \\ \rightarrow Z_{\text{tail}} \rightarrow b\bar{b}. \end{aligned} \quad (4)$$

In Fig. 3 we plot the Higgs resonance signals and various types of background for this mass range. While the cross sections are fairly large, the requirements on the beam energy resolution of the $\mu^+\mu^-$ collider are very constraining. In Fig. 4 we show the relationship between the Higgs width and the required machine energy resolution. If the resolution requirements can be met, the machine luminosity of $\sim 10^{31}$ $\text{cm}^{-2} \text{s}^{-1}$ could be adequate to facilitate the discovery of the Higgs in the mass range of 100–180 GeV.

For masses above 180 GeV, the dominant Higgs decay is

$$H^0 \rightarrow W^+W^- \quad \text{or} \quad Z^0Z^0 \quad (5)$$




Fig. 1. Schematic of a possible $\mu\mu$ collider scheme (few hundred GeV).




Fig. 2. Upper and lower bounds on m_{ϕ^0} as a function of m_t , coming from the requirement of a perturbative theory [4].

giving a clear signal and a larger width; the machine energy resolution requirements could be relaxed somewhat!

Another possibility for the intermediate Higgs mass range is to search for

$$\mu^+ \mu^- \rightarrow Z^0 + H^0 \quad (6)$$




Fig. 3. Estimated cross sections and backgrounds for Higgs production in the direct channel or by associated production for various Higgs mass.




Fig. 4. Higgs search at a $\mu^+\mu^-$ collider. Required machine resolution and the expected Higgs width.

using a broad energy sweep. The corresponding cross section is small (see Figs. 3 and 5). Once an approximate mass is determined, a strategy for the energy sweep through the resonance can be devised. The study of the t quark through $t\bar{t}$ production would also be interesting.

Finally, another possibility is to use the polarization of the $\mu^+\mu^-$ particles orientated so that only scalar interactions are possible (thus eliminating the background from single photon intermediate states as shown in Fig. 3) [6]. However, there would be a trade-off with luminosity and thus a strategy would have to be devised to maximize the possibility of success in the energy sweep through the resonance.

At the Napa workshop the possibility of developing a $\mu^+\mu^-$ collider in the $10^{31} \text{ cm}^{-2} \text{ s}^{-1}$ region was considered and appears feasible. It is less certain that the high energy resolution required for the Higgs sweep can be




Fig. 5. Total cross section for $\mu^+\mu^- \rightarrow \phi^0 Z$ as a function of \sqrt{s} for the fixed m_{ϕ_0} values indicated by the numbers (in GeV) beside each line. (Adapted from Ref. [4].)

Table 1
Key issues in the Higgs search

There is growing evidence that one Higgs particle is below $2 M_Z$ SUSY Higgs – 3 Higgs – one near M_Z (possibly up to ~ 130 GeV); extremely hard to detect

Hadron machines can search for these Higgs provided:

- i $\int \mathcal{L} dt \geq 10^5 \text{ pb}$ (LHC);
- ii the background for $H \rightarrow \gamma\gamma$ is small enough

A $\mu^+\mu^-$ collider with $\mathcal{L} \geq 10^{31} \text{ cm}^{-2} \text{ s}^{-1}$ operating between 100–180 GeV could discover the Higgs in 2000+ provided sufficient energy resolution is achieved

obtained. We summarize in Table 1 some of the key issues in the Higgs search.

$\mu^+\mu^-$ colliders could also be very important in the TeV energy range; however, since the cross sections for new particle production are much smaller, the luminosity requirements would be $\mathcal{L}_{\mu^+\mu^-} \geq 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$. This is the energy range where e^+e^- linear colliders are extremely difficult to develop [1]. This possibility will be the subject of a second Napa workshop to be held in 1994.

Acknowledgement

I wish to thank the participants of the workshop, and in particular Dave Neuffer and Andy Sessler, Bill Barletta and Kirk McDonald for discussions.

References

- [1] M. Tigner, presented at the 3rd Int. Workshop on Advanced Accelerator Concepts, Port Jefferson, NY, (1992); other paper to be published in AIP, 1992, ed. J. Wurtle.
- [2] Early references for $\mu\mu$ colliders are: E.A. Perevedentsev and A.N. Skrinsky, Proc. 12th Int. Conf. on High Energy Accelerators eds. F.T. Cole and R. Donaldson, 1983, p. 485; D. Neuffer, Proc. 12th Int. Conf. on High Energy Accelerators, eds F.T. Cole and R. Donaldson, 1983, p. 481; D. Neuffer, Particle Accel. 14 (1983) 75; D. Neuffer, in: Advanced Accelerator Concepts, AIP Conf. Proc. 156 (1987) 201.
- [3] Reports from the Napa Workshop: D. Neuffer, $\mu^+\mu^-$ Colliders: Possibilities and Challenges; W. Barletta and A. Sessler, Characteristics of a high energy $\mu^+\mu^-$ collider based on electro-production of muons.
- [4] See for primary references to the theoretical estimates here S. Dawson, J.F. Gunion, H.E. Haber and G.L. Kane, The Physics of the Higgs Bosons: Higgs Hunter's Guide (Addison Wesley, Menlo Park, 1989).
- [5] For example the Compact Muon Solenoid detector at the LHC, CERN reports.
- [6] This possibility came up in discussion with K. McDonald, Princeton.