

Measurements of the Branching Fractions of Exclusive Charmless B Meson Decays with η' or ω Mesons

B. Aubert,¹ D. Boutigny,¹ J.-M. Gaillard,¹ A. Hicheur,¹ Y. Karyotakis,¹ J.P. Lees,¹ P. Robbe,¹ V. Tisserand,¹ A. Palano,² G.P. Chen,³ J.C. Chen,³ N.D. Qi,³ G. Rong,³ P. Wang,³ Y.S. Zhu,³ G. Eigen,⁴ P.L. Reinertsen,⁴ B. Stugu,⁴ B. Abbott,⁵ G.S. Abrams,⁵ A.W. Borgland,⁵ A.B. Breon,⁵ D.N. Brown,⁵ J. Button-Shafer,⁵ R.N. Cahn,⁵ A.R. Clark,⁵ M.S. Gill,⁵ A. Gritsan,⁵ Y. Groyzman,⁵ R.G. Jacobsen,⁵ R.W. Kadel,⁵ J. Kadyk,⁵ L.T. Kerth,⁵ S. Kluth,⁵ Yu. G. Kolomensky,⁵ J.F. Kral,⁵ C. LeClerc,⁵ M.E. Levi,⁵ T. Liu,⁵ G. Lynch,⁵ A.B. Meyer,⁵ M. Momayezi,⁵ P.J. Oddone,⁵ A. Perazzo,⁵ M. Pripstein,⁵ N.A. Roe,⁵ A. Romosan,⁵ M.T. Ronan,⁵ V.G. Shelkov,⁵ A.V. Telnov,⁵ W.A. Wenzel,⁵ P.G. Bright-Thomas,⁶ T.J. Harrison,⁶ C.M. Hawkes,⁶ D.J. Knowles,⁶ S.W. O'Neale,⁶ R.C. Penny,⁶ A.T. Watson,⁶ N.K. Watson,⁶ T. Deppermann,⁷ K. Goetzen,⁷ H. Koch,⁷ J. Krug,⁷ M. Kunze,⁷ B. Lewandowski,⁷ K. Peters,⁷ H. Schmuecker,⁷ M. Steinke,⁷ J.C. Andres,⁸ N.R. Barlow,⁸ W. Bhimji,⁸ N. Chevalier,⁸ P.J. Clark,⁸ W.N. Cottingham,⁸ N. De Groot,⁸ N. Dyce,⁸ B. Foster,⁸ J.D. McFall,⁸ D. Wallom,⁸ F.F. Wilson,⁸ K. Abe,⁹ C. Hearty,⁹ T.S. Mattison,⁹ J.A. McKenna,⁹ D. Thiessen,⁹ S. Jolly,¹⁰ A.K. McKemey,¹⁰ J. Tinslay,¹⁰ V.E. Blinov,¹¹ A.D. Bukin,¹¹ D.A. Bukin,¹¹ A.R. Buzykaev,¹¹ V.B. Golubev,¹¹ V.N. Ivanchenko,¹¹ A.A. Korol,¹¹ E.A. Kravchenko,¹¹ A.P. Onuchin,¹¹ A.A. Salnikov,¹¹ S.I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ V.I. Telnov,¹¹ A.N. Yushkov,¹¹ D. Best,¹² A.J. Lankford,¹² M. Mandelkern,¹² S. McMahon,¹² D.P. Stoker,¹² A. Ahsan,¹³ K. Arisaka,¹³ C. Buchanan,¹³ S. Chun,¹³ J.G. Branson,¹⁴ D.B. MacFarlane,¹⁴ S. Prell,¹⁴ Sh. Rahatlou,¹⁴ G. Raven,¹⁴ V. Sharma,¹⁴ C. Campagnari,¹⁵ B. Dahmes,¹⁵ P.A. Hart,¹⁵ N. Kuznetsova,¹⁵ S.L. Levy,¹⁵ O. Long,¹⁵ A. Lu,¹⁵ J.D. Richman,¹⁵ W. Verkerke,¹⁵ M. Witherell,¹⁵ S. Yellin,¹⁵ J. Beringer,¹⁶ D.E. Dorfan,¹⁶ A.M. Eisner,¹⁶ A. Frey,¹⁶ A.A. Grillo,¹⁶ M. Grothe,¹⁶ C.A. Heusch,¹⁶ R.P. Johnson,¹⁶ W. Kroeger,¹⁶ W.S. Lockman,¹⁶ T. Pulliam,¹⁶ H. Sadrozinski,¹⁶ T. Schalk,¹⁶ R.E. Schmitz,¹⁶ B.A. Schumm,¹⁶ A. Seiden,¹⁶ M. Turri,¹⁶ W. Walkowiak,¹⁶ D.C. Williams,¹⁶ M.G. Wilson,¹⁶ E. Chen,¹⁷ G.P. Dubois-Felsmann,¹⁷ A. Dvoretzskii,¹⁷ D.G. Hitlin,¹⁷ S. Metzler,¹⁷ J. Oyang,¹⁷ F.C. Porter,¹⁷ A. Ryd,¹⁷ A. Samuel,¹⁷ M. Weaver,¹⁷ S. Yang,¹⁷ R.Y. Zhu,¹⁷ S. Devmal,¹⁸ T.L. Geld,¹⁸ S. Jayatilleke,¹⁸ G. Mancinelli,¹⁸ B.T. Meadows,¹⁸ M.D. Sokoloff,¹⁸ T. Barillari,¹⁹ P. Bloom,¹⁹ M.O. Dima,¹⁹ S. Fahey,¹⁹ W.T. Ford,¹⁹ T.L. Hall,¹⁹ D.R. Johnson,¹⁹ U. Nauenberg,¹⁹ A. Olivas,¹⁹ H. Park,¹⁹ P. Rankin,¹⁹ J. Roy,¹⁹ S. Sen,¹⁹ J.G. Smith,¹⁹ W.C. van Hoek,¹⁹ D.L. Wagner,¹⁹ J. Blouw,²⁰ J.L. Harton,²⁰ M. Krishnamurthy,²⁰ A. Soffer,²⁰ W.H. Toki,²⁰ R.J. Wilson,²⁰ J. Zhang,²⁰ T. Brandt,²¹ J. Brose,²¹ T. Colberg,²¹ G. Dahlinger,²¹ M. Dickopp,²¹ R.S. Dubitzky,²¹ E. Maly,²¹ R. Müller-Pfefferkorn,²¹ S. Otto,²¹ K.R. Schubert,²¹ R. Schwierz,²¹ B. Spaan,²¹ L. Wilden,²¹ L. Behr,²² D. Bernard,²² G.R. Bonneaud,²² F. Brochard,²² J. Cohen-Tanugi,²² S. Ferrag,²² E. Roussot,²² S. T'Jampens,²² C. Thiebaux,²² G. Vasileiadis,²² M. Verderi,²² A. Anjomshoaa,²³ R. Bernet,²³ A. Khan,²³ F. Muheim,²³ S. Playfer,²³ J.E. Swain,²³ M. Falbo,²⁴ C. Borean,²⁵ C. Bozzi,²⁵ S. Dittongo,²⁵ M. Folegani,²⁵ L. Piemontese,²⁵ E. Treadwell,²⁶ F. Anulli,^{27,*} R. Baldini-Ferrolì,²⁷ A. Calcaterra,²⁷ R. de Sangro,²⁷ D. Falciai,²⁷ G. Finocchiaro,²⁷ P. Patteri,²⁷ I.M. Peruzzi,^{27,*} M. Piccolo,²⁷ Y. Xie,²⁷ A. Zallo,²⁷ S. Bagnasco,²⁸ A. Buzzo,²⁸ R. Contri,²⁸ G. Crosetti,²⁸ P. Fabbriatore,²⁸ S. Farinon,²⁸ M. Lo Vetere,²⁸ M. Macri,²⁸ M.R. Monge,²⁸ R. Musenich,²⁸ M. Pallavicini,²⁸ R. Parodi,²⁸ S. Passaggio,²⁸ F.C. Pastore,²⁸ C. Patrignani,²⁸ M.G. Pia,²⁸ C. Priano,²⁸ E. Robutti,²⁸ A. Santroni,²⁸ M. Morii,²⁹ R. Bartoldus,³⁰ T. Dignan,³⁰ R. Hamilton,³⁰ U. Mallik,³⁰ J. Cochran,³¹ H.B. Crawley,³¹ P.-A. Fischer,³¹ J. Lamsa,³¹ W.T. Meyer,³¹ E.I. Rosenberg,³¹ M. Benkebil,³² G. Grosdidier,³² C. Hast,³² A. Höcker,³² H.M. Lacker,³² V. LePeltier,³² A.M. Lutz,³² S. Plaszczynski,³² M.H. Schune,³² S. Trincaz-Duvoid,³² A. Valassi,³² G. Wormser,³² R.M. Bionta,³³ V. Brigljević,³³ D.J. Lange,³³ M. Mugge,³³ X. Shi,³³ K. van Bibber,³³ T.J. Wenaus,³³ D.M. Wright,³³ C.R. Wuest,³³ M. Carroll,³⁴ J.R. Fry,³⁴ E. Gabathuler,³⁴ R. Gamet,³⁴ M. George,³⁴ M. Kay,³⁴ D.J. Payne,³⁴ R.J. Sloane,³⁴ C. Touramanis,³⁴ M.L. Aspinwall,³⁵ D.A. Bowerman,³⁵ P.D. Dauncey,³⁵ U. Egede,³⁵ I. Eschrich,³⁵ N.J.W. Gunawardane,³⁵ J.A. Nash,³⁵ P. Sanders,³⁵ D. Smith,³⁵ D.E. Azzopardi,³⁶ J.J. Back,³⁶ P. Dixon,³⁶ P.F. Harrison,³⁶ R.J.L. Potter,³⁶ H.W. Shorthouse,³⁶ P. Strother,³⁶ P.B. Vidal,³⁶ M.I. Williams,³⁶ G. Cowan,³⁷ S. George,³⁷ M.G. Green,³⁷ A. Kurup,³⁷ C.E. Marker,³⁷ P. McGrath,³⁷ T.R. McMahon,³⁷ S. Ricciardi,³⁷ F. Salvatore,³⁷ I. Scott,³⁷ G. Vaitsas,³⁷ D. Brown,³⁸ C.L. Davis,³⁸ J. Allison,³⁹ R.J. Barlow,³⁹ J.T. Boyd,³⁹ A.C. Forti,³⁹ J. Fullwood,³⁹ F. Jackson,³⁹ G.D. Lafferty,³⁹ N. Savvas,³⁹ E.T. Simopoulos,³⁹ J.H. Weatherall,³⁹ A. Farbin,⁴⁰ A. Jawahery,⁴⁰ V. Lillard,⁴⁰ J. Olsen,⁴⁰ D.A. Roberts,⁴⁰ J.R. Schieck,⁴⁰ G. Blaylock,⁴¹ C. Dallapiccola,⁴¹ K.T. Flood,⁴¹ S.S. Hertzbach,⁴¹ R. Kofler,⁴¹ T.B. Moore,⁴¹ H. Staengle,⁴¹ S. Willocq,⁴¹ B. Brau,⁴² R. Cowan,⁴² G. Sciolla,⁴² F. Taylor,⁴² R.K. Yamamoto,⁴² M. Milek,⁴³ P.M. Patel,⁴³ J. Trischuk,⁴³ F. Lanni,⁴⁴ F. Palombo,⁴⁴ J.M. Bauer,⁴⁵ M. Booke,⁴⁵ L. Cremaldi,⁴⁵ V. Eschenburg,⁴⁵ R. Kroeger,⁴⁵ J. Reidy,⁴⁵ D.A. Sanders,⁴⁵ D.J. Summers,⁴⁵ J.P. Martin,⁴⁶

J. Y. Nief,⁴⁶ R. Seitz,⁴⁶ P. Taras,⁴⁶ V. Zacek,⁴⁶ H. Nicholson,⁴⁷ C. S. Sutton,⁴⁷ C. Cartaro,⁴⁸ N. Cavallo,^{48,†}
 G. De Nardo,⁴⁸ F. Fabozzi,⁴⁸ C. Gatto,⁴⁸ L. Lista,⁴⁸ P. Paolucci,⁴⁸ D. Piccolo,⁴⁸ C. Sciacca,⁴⁸ J. M. LoSecco,⁴⁹
 J. R. G. Alsmiller,⁵⁰ T. A. Gabriel,⁵⁰ T. Handler,⁵⁰ J. Brau,⁵¹ R. Frey,⁵¹ M. Iwasaki,⁵¹ N. B. Sinev,⁵¹ D. Strom,⁵¹
 F. Colecchia,⁵² F. Dal Corso,⁵² A. Dorigo,⁵² F. Galeazzi,⁵² M. Margoni,⁵² G. Michelon,⁵² M. Morandin,⁵² M. Posocco,⁵²
 M. Rotondo,⁵² F. Simonetto,⁵² R. Stroili,⁵² E. Torassa,⁵² C. Voci,⁵² M. Benayoun,⁵³ H. Briand,⁵³ J. Chauveau,⁵³
 P. David,⁵³ C. De la Vaissière,⁵³ L. Del Buono,⁵³ O. Hamon,⁵³ F. Le Diberder,⁵³ Ph. Leruste,⁵³ J. Lory,⁵³ L. Roos,⁵³
 J. Stark,⁵³ S. Versillé,⁵³ P. F. Manfredi,⁵⁴ V. Re,⁵⁴ V. Speziali,⁵⁴ E. D. Frank,⁵⁵ L. Gladney,⁵⁵ Q. H. Guo,⁵⁵ J. H. Panetta,⁵⁵
 C. Angelini,⁵⁶ G. Batignani,⁵⁶ S. Bettarini,⁵⁶ M. Bondioli,⁵⁶ M. Carpinelli,⁵⁶ F. Forti,⁵⁶ M. A. Giorgi,⁵⁶ A. Lusiani,⁵⁶
 F. Martinez-Vidal,⁵⁶ M. Morganti,⁵⁶ N. Neri,⁵⁶ E. Paoloni,⁵⁶ M. Rama,⁵⁶ G. Rizzo,⁵⁶ F. Sandrelli,⁵⁶ G. Simi,⁵⁶
 G. Triggiani,⁵⁶ J. Walsh,⁵⁶ M. Haire,⁵⁷ D. Judd,⁵⁷ K. Paick,⁵⁷ L. Turnbull,⁵⁷ D. E. Wagoner,⁵⁷ J. Albert,⁵⁸ C. Bula,⁵⁸
 P. Elmer,⁵⁸ C. Lu,⁵⁸ K. T. McDonald,⁵⁸ V. Miftakov,⁵⁸ S. F. Schaffner,⁵⁸ A. J. S. Smith,⁵⁸ A. Tumanov,⁵⁸ E. W. Varnes,⁵⁸
 G. Cavoto,⁵⁹ D. del Re,⁵⁹ R. Faccini,^{14,59} F. Ferrarotto,⁵⁹ F. Ferroni,⁵⁹ K. Fratini,⁵⁹ E. Lamanna,⁵⁹ E. Leonardi,⁵⁹
 M. A. Mazzoni,⁵⁹ S. Morganti,⁵⁹ G. Piredda,⁵⁹ F. Safai Tehrani,⁵⁹ M. Serra,⁵⁹ C. Voena,⁵⁹ S. Christ,⁶⁰ R. Waldi,⁶⁰
 T. Adye,⁶¹ B. Franek,⁶¹ N. I. Geddes,⁶¹ G. P. Gopal,⁶¹ S. M. Xella,⁶¹ R. Aleksan,⁶² G. De Domenico,⁶² S. Emery,⁶²
 A. Gaidot,⁶² S. F. Ganzhur,⁶² P.-F. Giraud,⁶² G. Hamel de Monchenault,⁶² W. Kozanecki,⁶² M. Langer,⁶²
 G. W. London,⁶² B. Mayer,⁶² B. Serfass,⁶² G. Vasseur,⁶² C. Yeche,⁶² M. Zito,⁶² N. Copty,⁶³ M. V. Purohit,⁶³
 H. Singh,⁶³ F. X. Yumiceva,⁶³ I. Adam,⁶⁴ P. L. Anthony,⁶⁴ D. Aston,⁶⁴ K. Baird,⁶⁴ E. Bloom,⁶⁴ A. M. Boyarski,⁶⁴
 F. Bulos,⁶⁴ G. Calderini,⁶⁴ R. Claus,⁶⁴ M. R. Convery,⁶⁴ D. P. Coupal,⁶⁴ D. H. Coward,⁶⁴ J. Dorfan,⁶⁴ M. Doser,⁶⁴
 W. Dunwoodie,⁶⁴ R. C. Field,⁶⁴ T. Glanzman,⁶⁴ G. L. Godfrey,⁶⁴ S. J. Gowdy,⁶⁴ P. Grosso,⁶⁴ T. Himel,⁶⁴
 M. E. Huffer,⁶⁴ W. R. Innes,⁶⁴ C. P. Jessop,⁶⁴ M. H. Kelsey,⁶⁴ P. Kim,⁶⁴ M. L. Kocian,⁶⁴ U. Langenegger,⁶⁴
 D. W. G. S. Leith,⁶⁴ S. Luitz,⁶⁴ V. Luth,⁶⁴ H. L. Lynch,⁶⁴ H. Marsiske,⁶⁴ S. Menke,⁶⁴ R. Messner,⁶⁴ K. C. Moffeit,⁶⁴
 R. Mount,⁶⁴ D. R. Muller,⁶⁴ C. P. O'Grady,⁶⁴ M. Perl,⁶⁴ S. Petrak,⁶⁴ H. Quinn,⁶⁴ B. N. Ratcliff,⁶⁴ S. H. Robertson,⁶⁴
 L. S. Rochester,⁶⁴ A. Roodman,⁶⁴ T. Schietinger,⁶⁴ R. H. Schindler,⁶⁴ J. Schwiening,⁶⁴ V. V. Serbo,⁶⁴ A. Snyder,⁶⁴
 A. Soha,⁶⁴ S. M. Spanier,⁶⁴ J. Stelzer,⁶⁴ D. Su,⁶⁴ M. K. Sullivan,⁶⁴ H. A. Tanaka,⁶⁴ J. Va'vra,⁶⁴ S. R. Wagner,⁶⁴
 A. J. R. Weinstein,⁶⁴ W. J. Wisniewski,⁶⁴ D. H. Wright,⁶⁴ C. C. Young,⁶⁴ P. R. Burchat,⁶⁵ C. H. Cheng,⁶⁵ D. Kirkby,⁶⁵
 T. I. Meyer,⁶⁵ C. Roat,⁶⁵ R. Henderson,⁶⁶ W. Bugg,⁶⁷ H. Cohn,⁶⁷ A. W. Weidemann,⁶⁷ J. M. Izen,⁶⁸ I. Kitayama,⁶⁸
 X. C. Lou,⁶⁸ M. Turcotte,⁶⁸ F. Bianchi,⁶⁹ M. Bona,⁶⁹ B. Di Girolamo,⁶⁹ D. Gamba,⁶⁹ A. Smol,⁶⁹ D. Zanin,⁶⁹
 L. Lanceri,⁷⁰ A. Pompili,⁷⁰ G. Vuagnin,⁷⁰ R. S. Panvini,⁷¹ C. M. Brown,⁷² A. De Silva,⁷² R. Kowalewski,⁷²
 J. M. Roney,⁷² H. R. Band,⁷³ E. Charles,⁷³ S. Dasu,⁷³ F. Di Lodovico,⁷³ A. M. Eichenbaum,⁷³ H. Hu,⁷³ J. R. Johnson,⁷³
 R. Liu,⁷³ J. Nielsen,⁷³ Y. Pan,⁷³ R. Prepost,⁷³ I. J. Scott,⁷³ S. J. Sekula,⁷³ J. H. von Wimmersperg-Toeller,⁷³ S. L. Wu,⁷³
 Z. Yu,⁷³ H. Zoernig,⁷³ T. M. B. Kordich,⁷⁴ and H. Neal⁷⁴

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

³Institute of High Energy Physics, Beijing 100039, China

⁴University of Bergen, Institute of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720

⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁷Ruhr Universität Bochum, Institut für Experimentalphysik I, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 1TL, United Kingdom

⁹University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, California 92697

¹³University of California at Los Angeles, Los Angeles, California 90024

¹⁴University of California at San Diego, La Jolla, California 92093

¹⁵University of California at Santa Barbara, Santa Barbara, California 93106

¹⁶University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064

¹⁷California Institute of Technology, Pasadena, California 91125

¹⁸University of Cincinnati, Cincinnati, Ohio 45221

¹⁹University of Colorado, Boulder, Colorado 80309

²⁰Colorado State University, Fort Collins, Colorado 80523

²¹Technische Universität Dresden, Institut für Kern und Teilchenphysik, D-01062, Dresden, Germany

²²Ecole Polytechnique, F-91128 Palaiseau, France

²³University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

- ²⁴Elon College, Elon College, North Carolina 27244-2010
- ²⁵Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
- ²⁶Florida A&M University, Tallahassee, Florida 32307
- ²⁷Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
- ²⁸Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
- ²⁹Harvard University, Cambridge, Massachusetts 02138
- ³⁰University of Iowa, Iowa City, Iowa 52242
- ³¹Iowa State University, Ames, Iowa 50011-3160
- ³²Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
- ³³Lawrence Livermore National Laboratory, Livermore, California 94550
- ³⁴University of Liverpool, Liverpool L69 3BX, United Kingdom
- ³⁵University of London, Imperial College, London, SW7 2BW, United Kingdom
- ³⁶Queen Mary, University of London, E1 4NS, United Kingdom
- ³⁷University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
- ³⁸University of Louisville, Louisville, Kentucky 40292
- ³⁹University of Manchester, Manchester M13 9PL, United Kingdom
- ⁴⁰University of Maryland, College Park, Maryland 20742
- ⁴¹University of Massachusetts, Amherst, Massachusetts 01003
- ⁴²Massachusetts Institute of Technology, Lab for Nuclear Science, Cambridge, Massachusetts 02139
- ⁴³McGill University, Montréal, Canada QC H3A 2T8
- ⁴⁴Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
- ⁴⁵University of Mississippi, University, Mississippi 38677
- ⁴⁶Université de Montréal, Laboratoire Rene J.A. Levesque, Montréal, Canada QC H3C 3J7
- ⁴⁷Mount Holyoke College, South Hadley, Massachusetts 01075
- ⁴⁸Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
- ⁴⁹University of Notre Dame, Notre Dame, Indiana 46556
- ⁵⁰Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831
- ⁵¹University of Oregon, Eugene, Oregon 97403
- ⁵²Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
- ⁵³Universités Paris VI et VII, LPNHE, F-75252 Paris, France
- ⁵⁴Università di Pavia, Dipartimento di Eletttronica and INFN, I-27100 Pavia, Italy
- ⁵⁵University of Pennsylvania, Philadelphia, Pennsylvania 19104
- ⁵⁶Università di Pisa, Scuola Normale Superiore and INFN, I-56010 Pisa, Italy
- ⁵⁷Prairie View A&M University, Prairie View, Texas 77446
- ⁵⁸Princeton University, Princeton, New Jersey 08544
- ⁵⁹Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
- ⁶⁰Universität Rostock, D-18051 Rostock, Germany
- ⁶¹Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
- ⁶²DAPNIA, Commissariat à l'Energie Atomique/Saclay, F-91191 Gif-sur-Yvette, France
- ⁶³University of South Carolina, Columbia, South Carolina 29208
- ⁶⁴Stanford Linear Accelerator Center, Stanford, California 94309
- ⁶⁵Stanford University, Stanford, California 94305-4060
- ⁶⁶TRIUMF, Vancouver, British Columbia Canada V6T 2A3
- ⁶⁷University of Tennessee, Knoxville, Tennessee 37996
- ⁶⁸University of Texas at Dallas, Richardson, Texas 75083
- ⁶⁹Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
- ⁷⁰Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
- ⁷¹Vanderbilt University, Nashville, Tennessee 37235
- ⁷²University of Victoria, Victoria, British Columbia Canada V8W 3P6
- ⁷³University of Wisconsin, Madison, Wisconsin 53706
- ⁷⁴Yale University, New Haven, Connecticut 06511

(Received 7 August 2001; published 12 November 2001)

We present the results of searches for B decays to charmless two-body final states containing η' or ω mesons, based on 20.7 fb^{-1} of data collected with the BABAR detector. We find the branching fractions $\mathcal{B}(B^+ \rightarrow \eta' K^+) = (70 \pm 8 \pm 5) \times 10^{-6}$, $\mathcal{B}(B^0 \rightarrow \eta' K^0) = (42_{-11}^{+13} \pm 4) \times 10^{-6}$, and $\mathcal{B}(B^+ \rightarrow \omega \pi^+) = (6.6_{-1.8}^{+2.1} \pm 0.7) \times 10^{-6}$, where the first error quoted is statistical and the second is systematic. We give measurements of four additional modes for which the 90% confidence level upper limits are $\mathcal{B}(B^+ \rightarrow \eta' \pi^+) < 12 \times 10^{-6}$, $\mathcal{B}(B^+ \rightarrow \omega K^+) < 4 \times 10^{-6}$, $\mathcal{B}(B^0 \rightarrow \omega K^0) < 13 \times 10^{-6}$, and $\mathcal{B}(B^0 \rightarrow \omega \pi^0) < 3 \times 10^{-6}$.

We report results of searches for B decays to the charmless two-body final states [1] $B^+ \rightarrow \omega\pi^+$, $B^+ \rightarrow \omega K^+$, $B^+ \rightarrow \eta'\pi^+$, $B^+ \rightarrow \eta'K^+$, $B^0 \rightarrow \omega K^0$, $B^0 \rightarrow \omega\pi^0$, and $B^0 \rightarrow \eta'K^0$. These processes are manifestations of penguin or suppressed tree amplitudes proportional to small couplings in hadronic flavor mixing [Cabibbo-Kobayashi-Maskawa (CKM) matrix [2]]. Because of the absence of CKM favored $b \rightarrow c$ amplitudes, these decays are particularly sensitive to potentially new contributions from interference effects and virtual particles in loops. Previous measurements [3] yielded an unexpectedly large rate for $B \rightarrow \eta'K$, motivating a number of new theoretical ideas. The precise measurement of these and additional rare B decay modes will enable a better understanding of the underlying decay mechanism, including the possible contribution of physics beyond the standard model. This in turn will contribute to the measurement of fundamental parameters, including the CP -violating CKM phases.

The data were collected with the *BABAR* detector [4] at the PEP-II asymmetric e^+e^- collider [5] located at the Stanford Linear Accelerator Center. The results presented in this paper are based on data taken in the 1999–2000 run. An integrated luminosity of 20.7 fb^{-1} , corresponding to 22.7 million $B\bar{B}$ pairs, was recorded at the $Y(4S)$ resonance (“on resonance,” 10.58 GeV), with an additional 2.6 fb^{-1} about 40 MeV below this energy (“off resonance”) for the study of continuum backgrounds.

The asymmetric beam configuration in the laboratory frame provides a boost to the $Y(4S)$ increasing the momentum range of the B meson decay products up to $4.3 \text{ GeV}/c$. Charged particles are detected and their momenta are measured by a combination of a silicon vertex tracker, consisting of five layers of double-sided detectors, and a 40-layer central drift chamber, both operating in the 1.5 T magnetic field of a solenoid. Photons and electrons are detected by a CsI(Tl) electromagnetic calorimeter (EMC), which provides excellent angular and energy resolution with high efficiency for energies above 20 MeV [4].

Charged particle identification (PID) is provided by the average energy loss (dE/dx) in the tracking devices and by a unique, ring imaging detector of internally reflecting Cherenkov (DIRC) light covering the central region. A Cherenkov angle $K - \pi$ separation of better than four standard deviations (σ) is achieved for tracks below $3 \text{ GeV}/c$ momentum, decreasing to 2.5σ at the highest momenta in the final states considered here [6]. Electrons are identified with the use of the EMC.

We reconstruct a B meson candidate by combining an ω or η' candidate with a charged track, $\pi^0 \rightarrow \gamma\gamma$, or $K_S^0 \rightarrow \pi^+\pi^-$. The resonance decays R we reconstruct are $\omega \rightarrow \pi^+\pi^-\pi^0$, $\eta' \rightarrow \eta\pi^+\pi^-$ ($\eta'_\eta\pi\pi$), or $\eta' \rightarrow \rho^0\gamma$ ($\eta'_{\rho\gamma}$), with $\eta \rightarrow \gamma\gamma$ and $\rho^0 \rightarrow \pi^+\pi^-$. These modes are kinematically distinct from the dominant B decays to heavier charmed states. Backgrounds come primarily from combinatorics among continuum events in which a light quark pair is produced instead of an $Y(4S)$.

Monte Carlo (MC) simulations [7] of the target decay modes and of continuum background are used to establish the event selection criteria. The selection is designed to achieve high efficiency and retain sidebands sufficient to characterize the background for subsequent fitting. Photons must satisfy $E_\gamma > 50$ (100) MeV for π^0 (η) candidates. For $\eta' \rightarrow \rho^0\gamma$ candidates from $B^+ \rightarrow \eta'K^+$ and $B^+ \rightarrow \eta'\pi^+$ the requirement is $E_\gamma > 200$ MeV, while for $B^0 \rightarrow \eta'K^0$ it is looser ($E_\gamma > 100$ MeV) because of smaller combinatoric background.

We select ω , η' , η , and ρ candidates with the following requirements on the invariant masses in MeV/c^2 of their final states: $735 < m(\pi^+\pi^-\pi^0) < 830$, $930 < m(\eta\pi^+\pi^-) < 990$, $900 < m(\rho\gamma) < 1000$, $490 < m(\gamma\gamma) < 600$, and $500 < m(\pi^+\pi^-) < 995$. For π^0 and K_S^0 candidates we require $120 < m(\gamma\gamma) < 150$ and $488 < m(\pi^+\pi^-) < 508$.

Tracks in ω , η' , or ρ candidates must have DIRC, dE/dx , and EMC responses consistent with pions. For charged B decays, the B primary track must have an associated DIRC Cherenkov angle within 3.5σ of the expected value for a kaon or pion. For modes with K_S^0 the three-dimensional flight distance from the production point must exceed 2 mm, and the angle between the flight and momentum vectors projected perpendicular to the beam must be less than 40 mrad.

A B meson candidate is characterized by two kinematic observables. The minimally correlated pair we use is the energy constrained mass m_{EC} and energy difference ΔE . In the $Y(4S)$ frame the B meson energy E^* equals the beam energy E_{beam}^* . A kinematic fit of the measured candidate four momentum in this frame with the constraint $E^* = E_{\text{beam}}^*$ yields m_{EC} , while $\Delta E \equiv E^* - E_{\text{beam}}^*$ measures the consistency of this constraint. We require $|\Delta E| \leq 0.2 \text{ GeV}$ and $m_{EC} \geq 5.2 \text{ GeV}/c^2$. The resolutions on these quantities are mode dependent but average about 30 MeV and $2.8 \text{ MeV}/c^2$, respectively.

To discriminate against tau-pair and two-photon background we require the event to contain at least five charged tracks. To reject continuum background we make use of the angle θ_T between the thrust axes of the B candidate and the rest of the tracks and neutral clusters in the event, calculated in the center-of-mass frame. The distribution of $\cos\theta_T$ is sharply peaked near ± 1 for combinations drawn from jetlike $q\bar{q}$ pairs, and nearly uniform for the isotropic B meson decays.

The yields are obtained from extended unbinned maximum likelihood (ML) fits, with two variants specified in the following paragraphs. The first (ML1), which provides our results for all modes except $B^0 \rightarrow \omega\pi^0$, uses several uncorrelated variables for the kinematics of the B decay chain and a Fisher discriminant for the production and energy flow. The second (ML2) is applied to all channels with an ω meson; it uses ΔE and the output of a neural network built from the remaining inputs. Comparisons of the $\omega\pi^+$, ωK^+ , and ωK^0 modes show that the central values and errors for the yields obtained by the two approaches

are in very good agreement. Simple cut-based analyses are performed as checks for each final state. Agreement of central values is good in all cases, although, as expected, errors are larger than for the ML analyses, particularly for modes having high background.

The ML1 fit method is applied to events satisfying $|\cos\theta_T| \leq 0.9$. The input observables are ΔE , m_{EC} , the invariant mass m_R of the intermediate resonance, the Fisher discriminant \mathcal{F} , and, where relevant, the η mass m_η , the measured DIRC angle for the B primary track, and the cosine \mathcal{H} of the helicity angle, the angle in the ω rest frame between the normal to the ω decay plane and the B flight direction. The Fisher discriminant [8] combines eleven variables: the angles with respect to the beam axis in the $Y(4S)$ frame of the B momentum and the B two-body decay axis, and a nine bin representation of the energy flow about the B decay axis.

For the ML2 method we relax the preliminary requirements to $100 < m(\gamma\gamma) < 160$ MeV/ c^2 and $|\Delta E| < 0.3$ GeV. The neural network is constructed with the B momentum p^* , a χ^2 for resonance masses, \mathcal{H} , and variables representing energy flow and angular distributions, including θ_T .

We use MC to estimate backgrounds from other B decays, including final states with and without charm. For most of our modes we find contributions that are negligible. For the $\eta' \rightarrow \rho^0 \gamma$ modes we account for small cross feed contributions in the systematic error estimate.

The likelihood function for N events is

$$\mathcal{L} = \frac{e^{-(\sum n_j)}}{N!} \prod_{i=1}^N \mathcal{L}_i, \quad \mathcal{L}_i = \sum_{j=1}^m n_j \mathcal{P}_j(\mathbf{x}_i).$$

Here n_j is the population size for species j (e.g., signal, background) and $\mathcal{P}_j(\mathbf{x}_i)$ is the corresponding probability distribution function (PDF), evaluated with the observables \mathbf{x}_i of the i th event.

For the fits of charged B decays, \mathcal{L}_i becomes

$$\mathcal{L}_i = n_\pi \mathcal{P}_{\pi S}(\mathbf{x}_i) + n_K \mathcal{P}_{KS}(\mathbf{x}_i) + n_C [f_{KC} \mathcal{P}_{KC}(\mathbf{x}_i) + (1 - f_{KC}) \mathcal{P}_{\pi C}(\mathbf{x}_i)],$$

where n_π (n_K) is the number of $B^+ \rightarrow R\pi^+$ ($B^+ \rightarrow RK^+$) signal events, n_C is the number of continuum background events, and f_{KC} is the fraction of continuum background events for which the B primary track is identified as a kaon. These quantities are the free parameters of the ML fit. The probabilities for the components are $\mathcal{P}_{\pi S}$ (\mathcal{P}_{KS}) for $B^+ \rightarrow R\pi^+$ ($B^+ \rightarrow RK^+$) signal and $\mathcal{P}_{\pi C}$ (\mathcal{P}_{KC}) for background where the primary track is a pion (kaon). Since we measure the correlations among the observables in the data to be small, we take each \mathcal{P}_j to be a product of the PDFs for the separate observables. The analyses involving a K_S^0 are treated identically, except that there is only one component of signal and of continuum background.

A second B candidate satisfying the preliminary cuts occurs in about 10%–20% of the events. In this case the

“best” combination is selected according to a χ^2 quantity computed with m_{EC} , m_R , m_η (for $\eta' \rightarrow \eta\pi^+\pi^-$ modes), and the Fisher discriminant.

We determine the PDFs for the likelihood fits from simulation for the signal component, and from off-resonance and sideband data for the continuum background. Peaking distributions (signal masses, ΔE , \mathcal{F}) are parametrized as Gaussians, with or without a second Gaussian or asymmetric width as required to describe the distributions. Slowly varying distributions (combinatoric background under mass or energy peaks, \mathcal{H} or \mathcal{F}) have first or second order polynomial shapes. The combinatoric background in m_{EC} is described by a phase space motivated empirical function [9]. Control samples of B decays to charmed final states of similar topology are used to verify the simulated resolutions in ΔE and m_{EC} . Inclusive resonance production samples, such as those shown in Fig. 1, are used similarly for the relevant B daughter mass spectra.

We compute the branching fractions from the fitted signal event yields, reconstruction efficiency, daughter branching fractions, and the number of produced B mesons, assuming equal production rates of charged and neutral pairs. To determine the reconstruction efficiency, including any yield bias of the likelihood fit, we apply the method to simulated samples with the signal and continuum background populations expected in the data. Table I shows for each decay chain the branching fraction we measure, together with the quantities entering into its computation. The statistical error on the number of events is taken as the shift from the central value that changes the quantity $\chi^2 \equiv -2 \log(\mathcal{L} / \mathcal{L}_{\max})$ by one unit. We also give the significance S , computed as the square root of the difference between the value of χ^2 for zero signal and the value at its minimum. The χ^2 used for significance includes a term that accounts for the additive systematic




FIG. 1. Invariant mass distributions for inclusive data samples of candidates with $Y(4S)$ frame momentum greater than 2.3 GeV/ c for (a) η' , with $520 < m(\gamma\gamma) < 575$ MeV/ c^2 , and (b) ω candidates, with $120 < m(\gamma\gamma) < 150$ MeV/ c^2 . From the overlaid fit curves, the Gaussian peak widths are 4 and 10 MeV/ c^2 , respectively.

TABLE I. Signal event yield with statistical uncertainty, detection efficiency ϵ , daughter branching fractions that were forced to 100% in our signal mode simulations, significance S (defined in the text), and branching fraction result for each decay chain or mode, with the final (combined) result given in bold type. We show 90% confidence upper limits in parentheses where appropriate.

Mode	Yield	ϵ (%)	$\prod \mathcal{B}_i$ (%)	S (σ)	\mathcal{B} (10^{-6})
$\eta'_{\eta\pi\pi}K^+$	$49.5^{+8.1}_{-7.3}$	20	17.4	15	63^{+10}_{-9}
$\eta'_{\rho\gamma}K^+$	$87.6^{+13.4}_{-12.5}$	18	29.5	11	80^{+12}_{-11}
$\eta'K^+$	17	$70 \pm 8 \pm 5$
$\eta'_{\eta\pi\pi}K^0$	$6.3^{+3.3}_{-2.5}$	16	6.0	4.7	28^{+15}_{-11}
$\eta'_{\rho\gamma}K^0$	$20.8^{+7.4}_{-6.5}$	16	10.1	4.2	61^{+22}_{-19}
$\eta'K^0$	5.9	$42^{+13}_{-11} \pm 4$
$\eta'_{\eta\pi\pi}\pi^+$	$5.7^{+3.8}_{-2.8}$	20	17.4	3.2	$7.1^{+4.8}_{-3.5}$
$\eta'_{\rho\gamma}\pi^+$	$-0.9^{+7.8}_{-6.2}$	19	29.5	0.1	$-0.7^{+6.7}_{-5.3}$
$\eta'\pi^+$	2.8	$5.4^{+3.5}_{-2.6} \pm 0.8$ (<12)
ωK^+	$6.4^{+5.6}_{-4.4}$	22	88.8	1.3	$1.4^{+1.3}_{-1.0} \pm 0.3$ (<4)
ωK^0	$8.1^{+4.6}_{-3.6}$	18	30.5	3.2	$6.4^{+3.6}_{-2.8} \pm 0.8$ (<13)
$\omega\pi^+$	$27.6^{+8.8}_{-7.7}$	21	88.8	4.9	$6.6^{+2.1}_{-1.8} \pm 0.7$
$\omega\pi^0$	$-0.9^{+5.0}_{-3.2}$	18	88.8	...	$-0.3 \pm 1.1 \pm 0.3$ (<3)

error discussed below. Where the significance is less than four standard deviations, we quote also (Bayesian) 90% C.L. upper limits, defined by the solution B to the condition $\int_0^B \mathcal{L}(b) db / \int_0^\infty \mathcal{L}(b) db = 0.9$.

In Fig. 2 we show projections of m_{EC} and ΔE for the modes with significant yields. The projections are made by




FIG. 2. B candidate m_{EC} and ΔE for $B^+ \rightarrow \eta'K^+$ (a),(b), $B^0 \rightarrow \eta'K^0$ (c),(d), and $B^+ \rightarrow \omega\pi^+$ (e),(f). Histograms represent data, with the $\eta' \rightarrow \eta\pi\pi$ subset shaded, the solid curves represent the full fit functions, and the dashed curves represent the background functions.

selecting events with signal likelihood (computed without the variable plotted) exceeding a mode-dependent threshold that optimizes the expected sensitivity.

We have evaluated systematic errors, which are dominated in most cases by the PDF uncertainties (3%–18%, depending on the decay mode). To determine these we vary parameters of the PDFs within their uncertainties and observe the impact on the fit yield. We include them in upper limits by convolution with the likelihood function. This is the only additive systematic error; all others are multiplicative. The estimate of any systematic bias from the fitter itself (1%–4%) comes from fits of simulated samples with varying background populations.

The uncertainty in our knowledge of the efficiency is found from auxiliary studies to be 1% per track, 1.25% per photon, and 5% per K_S^0 for the candidate B and the unreconstructed B , which must contribute tracks to fulfill the event multiplicity requirement. We add these errors linearly for the required tracks in the event and, similarly, for the photons and neutral kaons. Our estimate of the B production systematic error is 1.6%. Published world averages [10] provide the B daughter branching fraction uncertainties.

Systematic errors associated with the event selection are minimal given the generally loose requirements. We account explicitly for $\cos\theta_T$ (1%), for which we observe a nearly uniform distribution in the signal simulation. We also include errors of 4% from those PID requirements that are imposed via cuts rather than the fit.

We observed signals of at least 4σ in five of the decay chains studied here, as reported in Table I. Where we have multiple chains for a given mode, we combine the results

by adding the χ^2 distributions that represent them and their uncorrelated statistical and systematic errors.

The final results are generally in agreement with those previously reported [3,11], with somewhat smaller errors. In particular, we confirm the expected $\mathcal{B}(B^+ \rightarrow \omega \pi^+) > \mathcal{B}(B^+ \rightarrow \omega K^+)$, and the rather larger than predicted [12] rate for $B \rightarrow \eta' K$ obtained by the CLEO Collaboration [3]. Conjectured sources of η' enhancement include flavor singlet [13], charm enhanced [14], and constructively interfering internal penguin diagrams [12,15]. Our results in combination with expected measurements of related modes involving η and K^* should help to clarify this situation.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (U.S.A.), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF (Germany), INFN (Italy), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from the Swiss NSF, A.P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

*Also with Università di Perugia, Perugia, Italy.

†Also with Università della Basilicata, Potenza, Italy.

[1] The charge conjugate of the named state is implicitly included, here and throughout this paper.

- [2] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [3] CLEO Collaboration, S.J. Richichi *et al.*, Phys. Rev. Lett. **85**, 520 (2000).
- [4] BABAR Collaboration, B. Aubert *et al.*, Report No. SLAC-PUB-8569, hep-ex/0105044 [Nucl. Instrum. Methods Phys. Res., Sect. A (to be published)].
- [5] PEP-II Conceptual Design Report No. SLAC-R-418, 1993.
- [6] See Fig. 1(b) in BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **87**, 151802 (2001).
- [7] The BABAR detector Monte Carlo simulation is based on GEANT; R. Brun *et al.*, CERN Report No. DD/EE/84-1.
- [8] CLEO Collaboration, D.M. Asner *et al.*, Phys. Rev. D **53**, 1039 (1996).
- [9] With $x \equiv m_{EC}/E_b$ and ξ a parameter to be fit, $f(x) \propto x\sqrt{1-x^2} \exp[-\xi(1-x^2)]$. See ARGUS Collaboration, H. Albrecht *et al.*, Phys. Lett. B **241**, 278 (1990).
- [10] Particle Data Group, D.E. Groom *et al.*, Eur. Phys. J. C **15**, 1 (2000).
- [11] CLEO Collaboration, C.P. Jessop *et al.*, Phys. Rev. Lett. **85**, 2881 (2000).
- [12] A. Ali, G. Kramer, and C.D. Lü, Phys. Rev. D **58**, 094009 (1998); Y.H. Chen *et al.*, Phys. Rev. D **60**, 094014 (1999).
- [13] M. Gronau and J.L. Rosner, Phys. Rev. D **53**, 2516 (1996); A.S. Dighe, M. Gronau, and J.L. Rosner, Phys. Rev. Lett. **79**, 4333 (1997); M.R. Ahmady, E. Kou, and A. Sugamoto, Phys. Rev. D **58**, 014015 (1998); D. Du, C.S. Kim, and Y. Yang, Phys. Lett. B **426**, 133 (1998).
- [14] I. Halperin and A. Zhitnitsky, Phys. Rev. D **56**, 7247 (1997); E.V. Shuryak and A. Zhitnitsky, *ibid.* **57**, 2001 (1998).
- [15] H.J. Lipkin, Phys. Lett. B **254**, 247 (1991).